

Can Spectroscopy Help Stem the Flow of Opioids?

Application Note



KEYWORDS

- Fentanyl
- Analogs
- Long fiber grating

TECHNIQUES

- Transmittance

APPLICATIONS

- Toxicology analysis
- Medical diagnostics

In this application, an HR series high-resolution spectrometer is used as part of a project to develop a fiber optic sensor that detects designer drugs, which mimic the effects of the powerful opioid fentanyl.

Background

The impact of the opioid crisis is staggering. According to the U.S. Department of Health and Human Services, more than 10 million people aged 12 or older abused opioids in 2019, and two-thirds of overdose-related deaths in the U.S. involved an opioid. Globally, 80% of premature death and disability related to substance abuse in otherwise healthy individuals has been attributed to opioids.¹

Now, the emergence of powerful synthetic opioids like fentanyl and its illicit analogs (chemically similar substances often referred to as designer drugs) has spurred demand for faster, less expensive detection methods. Could spectroscopy be part of an alternative to current toxicological analysis practices?

Solution

As described in a recent open-access paper published in *Biosensors and Bioelectronics*, a group of photonics and medical technology researchers from several European academic and business concerns has developed a novel fiber optic sensor for detecting fentanyl analogs in blood.²

US +1 727-733-2447

EUROPE +31 26-3190500

ASIA +86 21-6295-6600

info@oceanoptics.com • www.oceanoptics.com



OceanOptics

More specifically, the sensor is intended to detect butyrylfentanyl in blood. As of 2021, according to the paper's authors, regular toxicological analyses of this analog were not yet established.




Figure 1. A high-resolution spectrometer measures transmission of the sample in a flow system. An illustration of the long-period grating array is at top.

The research team's sensor consists of a long period fiber grating array with molecularly imprinted markers, with an Ocean Optics spectrometer to measure the transmission spectrum of the sensor for key analog criteria (**Figure 1**). Initial results have been successful, demonstrating how the sensor could provide an alternative to existing lab-based instruments that is faster, simpler, more compact and less expensive. What's more, the sensor would be adaptable for different opioids, helping law enforcement and medical professionals more readily keep pace with diagnostic challenges.

Equipment Used

- **HR series high-resolution spectrometer.** High-resolution spectrometers with <1.0 nm (FWHM) optical resolution that are ideal for applications including laser characterization, atomic emission lines analysis, and fiber grating transmission.
- **HL-2000 tungsten halogen light.** Provides stable output from 360-2400 nm.

Why Screening Matters

Although the opioid crisis to date has been largely confined to the U.S., interested parties in Europe and other regions have taken notice. That's partly because efforts to screen for new designer drugs are so challenging. For example, the U.S. Drug Enforcement Administration has identified nearly 20 different fentanyl analogs, some of which can be up to thousands of times more potent than morphine. Being able to rapidly and cost effectively screen for these illicit drugs using spectroscopy will help to save lives and reduce the strain on our health care resources already stretched because of the pandemic.

References

1. United Nations World Drug Report, 2020.
2. Liu, L., Grillo, F., Canfarotta, F., Whitcombe, M., Morgan, S.P., Piletsky, S., Correia, R., He, C., Norris, A., and Korposh, S. Carboxyl-fentanyl detection using optical fibre grating-based sensors functionalised with molecularly imprinted nanoparticles, *Biosensors and Bioelectronics*, Volume 177, 1 April 2021, 113002.